

KURZPROTOKOLL MIDOKIT

Öffentlicher Titel	Midostaurin und Standardchemotherapie bei KIT- oder FLT3-ITD positiver t(8;21) AML
Wissenschaftl. Titel	A single-arm phase II trial to assess the efficacy of midostaurin (PKC412) added to standard primary therapy in patients with newly diagnosed c-KIT or FLT3-ITD mutated t(8;21)AML
Kurztitel	MIDOKIT
Studienart	multizentrisch, prospektiv, offen/unverblindet, einarmig
Studienphase	Phase II
Erkrankung	Blut: Akute myeloische Leukämie (AML): Neu diagnostiziert / de novo
Ziele	<ul style="list-style-type: none">- To compare CR, OS, and CIR rates in the two arms- To compare the toxicity profile in the two arms- To compare the kinetics of MRD in the two arms- To compare the duration of patient hospitalization in the two arms- To compare the QoL between the two arms- To compare the event free survival rates in the two arms in the whole population
Einschlusskriterien	<ul style="list-style-type: none">- Signed written informed consent according to IGH/EU/GCP and national local laws- Newly diagnosed APL confirmed by the presence of t (15;17) or PML/RAR by RT-PCR or micro speckled PML nuclear distribution in leukemic cells- Age ≥ 18 and < 71 years- WHO performance status 0-2 included- Serum total bilirubin ≤ 3.0 mg/dl- Serum creatinine ≤ 3.0 mg/dl- WBC at diagnosis $\leq 10 \times 10^9 /L$
Ausschlusskriterien	<ul style="list-style-type: none">- Age < 18 and ≥ 71- WBC $> 10 \times 10^9 /L$- Active malignancy at time of study entry- Lack of diagnostic confirmation at genetic level- Significant arrhythmias, EKG abnormalities or neuropathy- Cardiac contraindications for intensive chemotherapy (L-EF $< 50\%$)- Uncontrolled, life-threatening infections- Severe uncontrolled pulmonary or cardiac disease- Women who are either pregnant or breast feeding, or of child-bearing potential, defined as all women physiologically capable of becoming pregnant, UNLESS they meet on eof the following definitions: Amenorrhea, post surgical bilateral oophorectomy with or without hysterectomy Using a highly effective method of birth control (defined as those which result in a failure rate less than 1 % per year) when used consistently an d correctly such as implants, injectables, oral contraceptives, IUDs, sexual abstinence or vasectomized partner- Concomitant severe psychiatric disorder- HIV positivity- Use of other investigational drugs at the time of enrolment or within 30 days before study entry
Alter	18 - 70 Jahre
Molekularer Marker	FLT3 AML-ETO KIT
Fallzahl	1

**KURZPROTOKOLL
MIDOKIT**

Prüfzentren

Universitätsklinikum Frankfurt (Rekrutierung beendet)
Medizinische Klinik II, Hämatologie/Onkologie
Theodor-Stern-Kai 7
60590 Frankfurt am Main
Sabine Hug
Tel: 069 6301-6353
Fax: 069 6301-7463
s.hug@em.uni-frankfurt.de

**Registrierung in anderen
Studienregistern**

ClinicalTrials.gov NCT01830361
EudraCT 2011-002567-17